


NCTN Leukemia Trials Portfolio (Open as of 11/15/2022)

Each far right box includes the NCTN protocol number with a hyperlink to the associated ClinicalTrials.gov webpage. Click on it to view the protocol title and study information.



NCTN Leukemia Trials (Open as of 11/15/2022)

| Protocol Number | Phase | Protocol Title |
|-----------------|--------|--|
| A041501 | III | A Phase III Trial to Evaluate the Efficacy of the Addition of Inotuzumab Ozogamicin (a Conjugated Anti-CD22 Monoclonal Antibody) to Frontline Therapy in Young Adults (Ages 18-39 Years) with Newly Diagnosed Precursor B-Cell ALL |
| A041701 | II/III | A Randomized Phase II/III Study of Conventional Chemotherapy +/- Uproleselan (GMI-1271) in Older Adults with Acute Myeloid Leukemia Receiving Intensive Induction Chemotherapy |
| A041703 | II | A Phase II Study of Inotuzumab Ozogamicin Followed by Blinatumomab for Ph-Negative, CD22-Positive B-Lineage Acute Lymphoblastic Leukemia in Newly Diagnosed Older Adults or Adults with Relapsed or Refractory Disease |
| EA9152 | I/II | A Phase IB/II Study of Venetoclax (ABT-199) in Combination with Liposomal Vincristine in Patients with Relapsed or Refractory T-Cell or B-Cell Acute Lymphoblastic Leukemia |
| EA9171 | II | Phase II Study of Adding the Anti-PD-1 Pembrolizumab to Tyrosine Kinase Inhibitors in Patients with Chronic Myeloid Leukemia and Persistently Detectable Minimal Residual Disease |
| EA9181 | III | A Phase III Randomized Trial of Steroids + Tyrosine Kinase Inhibitor Induction with Chemotherapy or Blinatumomab for Newly Diagnosed BCR-ABL-Positive Acute Lymphoblastic Leukemia in Adults |
| EA9213 | II | A Phase II Study of Daratumumab-Hyaluronidase for Chemotherapy-Relapsed/Refractory Minimal Residual Disease (MRD) in T Cell Acute Lymphoblastic Leukemia (T-ALL) |
| S1905 | II | A Phase I/II Study of AKR1C3-Activated Prodrug OBI-3424 (OBI-3424) In Patients with Relapsed/Refractory T-Cell Acute Lymphoblastic Leukemia (T-ALL)/T-Cell Lymphoblastic Lymphoma (T-LBL) |
| S1925 | III | Randomized, Phase III Study of Early Intervention with Venetoclax and Obinutuzumab Versus Delayed Therapy with Venetoclax and Obinutuzumab in Newly Diagnosed Asymptomatic High-Risk Patients with Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL): EVOLVE CLL/SLL Study |