


# NCTN Lymphoma Trials Portfolio (Open as of 9/15/2022)

Each far right box includes the NCTN protocol number with a hyperlink to the associated ClinicalTrials.gov webpage. Click on it to view the protocol title and study information.



## Cross-disease trial:

**EAY131 (MATCH)**

## NCTN Lymphoma Trials (Open as of /15/2022)

Protocol Number	Phase	Protocol Title
<b>A051301</b>	III	A Randomized Double-Blind Phase III Study of Ibrutinib During and Following Autologous Stem Cell Transplantation Versus Placebo in Patients with Relapsed or Refractory Diffuse Large B-cell Lymphoma of the Activated B-cell Subtype
<b>A051701</b>	II/III	Randomized Phase II/III Study of Venetoclax (ABT199) Plus Chemoimmunotherapy for MYC/BCL2 Double-Hit and Double Expressing Lymphomas
<b>A051901</b>	I	Phase I Trial of Methotrexate, Rituximab, Lenalidomide, and Nivolumab (Nivo-MR2) Induction Followed by Lenalidomide and Nivolumab Maintenance in Primary CNS Lymphoma
<b>A051902</b>	II	A Randomized Phase II Study of CHO(E)P vs CC-486-CHO(E)P vs Duvelisib-CHO(E)P in Previously Untreated CD30 Negative Peripheral T-Cell Lymphomas
<b>ANHL1931</b>	III	A Randomized Phase 3 Trial of Nivolumab (NSC# 748726) in Combination with Chemo-Immunotherapy for the Treatment of Newly Diagnosed Primary Mediastinal B-Cell Lymphoma
<b>E4412</b>	I/II	A Phase I Study with an Expansion Cohort/Randomized Phase II Study of the Combinations of Ipilimumab, Nivolumab and Brentuximab Vedotin in Patients with Relapsed/Refractory Hodgkin Lymphoma
<b>EA4151</b>	III	A Randomized Phase III Trial of Consolidation with Autologous Hematopoietic Cell Transplantation Followed by Maintenance Rituximab vs Maintenance Rituximab Alone for Patients with Mantle Cell Lymphoma in Minimal Residual Disease-Negative First Complete Remission
<b>EA4181</b>	II	A Randomized 3-Arm Phase II Study Comparing 1.) Bendamustine, Rituximab and High Dose Cytarabine (BR/CR) 2.) Bendamustine, Rituximab, High Dose Cytarabine and Acalabrutinib (BR/CR-A), and 3.) Bendamustine, Rituximab and Acalabrutinib (BR-A) in Patients <= 70 Years Old with Untreated Mantle Cell Lymphoma
<b>S1608</b>	II	Randomized Phase II Trial in Early Relapsing or Refractory Follicular Lymphoma
<b>S1826</b>	III	A Phase III, Randomized Study of Nivolumab (Opdivo) Plus AVD or Brentuximab Vedotin (Adcetris) Plus AVD in Patients (Age >= 12 Years) with Newly Diagnosed Advanced Stage Classical Hodgkin Lymphoma
<b>S1905</b>	II	A Phase I/II Study of AKR1C3-Activated Prodrug OBI-3424 (OBI-3424) In Patients with Relapsed/Refractory T-Cell Acute Lymphoblastic Leukemia (T-ALL)/T-Cell Lymphoblastic Lymphoma (T-LBL)
<b>S1918</b>	II/III	A Phase II/III Randomized Study of R-MiniCHOP with or Without CC-486 (Oral Azacitidine) in Participants Age 75 Years or Older with Newly Diagnosed Diffuse Large B Cell Lymphoma, Grade IIIB Follicular Lymphoma, Transformed Lymphoma, and High-Grade B-Cell Lymphomas with MYC AND BCL2 and/or BCL6 Rearrangements
<b>S1925</b>	III	Randomized, Phase III Study of Early Intervention with Venetoclax and Obinutuzumab Versus Delayed Therapy with Venetoclax and Obinutuzumab in Newly Diagnosed Asymptomatic High-Risk Patients with Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL): EVOLVE CLL/SLL Study
<b>S2005</b>	II	A Phase II Randomized Study of Ibrutinib and Rituximab with or Without Venetoclax in Previously Untreated Waldenström's Macroglobulinemia (WM) / Lymphoplasmacytic Lymphoma (LPL)
<b>EAY131</b>	II	Molecular Analysis for Therapy Choice (MATCH)